



## Crystal structure of seven-membered acetals with furan and pyridine planar fragments

V.Yu. Fedorenko<sup>a</sup>, O.A. Lodochnikova<sup>b</sup>, A.S. Petukhov<sup>a</sup>, O.N. Kataeva<sup>b</sup>, I.A. Litvinov<sup>b</sup>,  
Yu.G. Shtyrin<sup>a</sup>, E.N. Klimovitskii<sup>a,\*</sup>

<sup>a</sup>A.M. Butlerov Chemical Research Institute, Kazan State University, Kremlevskaya Str. 18, Kazan 420008, Russian Federation

<sup>b</sup>A.E. Arbusov Institute of Organic and Physical Chemistry, Russian Academy of Sciences, Arbuzov Str. 8, Kazan 420088, Russian Federation

Received 10 July 2002; revised 6 September 2002; accepted 6 September 2002

### Abstract

X-ray structure investigation of fused seven-membered acetals based on vitamin B<sub>6</sub> and 3,4-bis(hydroxymethyl)furan have been performed. Molecules adopt chair conformations with equatorial position of substituents at acetal carbons; the geometry of acetal cycles resembles that of related seven-membered phthalylacetals. Stereochemistry of the tetracyclic adduct of furan-containing acetal with maleic anhydride was also investigated. The product exhibits endo–exo configuration with appreciably distorted seven-membered chair-like conformation.

© 2002 Elsevier Science B.V. All rights reserved.

**Keywords:** Fused heterocycles; Seven-membered acetals; Furan and vitamin B<sub>6</sub> derivatives; X-ray crystallography

### 1. Introduction

Seven-membered cyclic acetals with a planar fragment (1,3-dioxacyclohept-5-ene and its derivatives) provide examples on the coexistence of chair and twist-boat forms in solutions [1–12]. Conformational equilibrium position depends on the volume of alkyl or aryl substituents at the acetal carbon atom and the type of the planar fragment [2,8]. The Taft postulate was drawn on the series of conformationally inhomogeneous 2-R-1,3-dioxacyclohept-5-enes [13] in bromination [14] and [4 + 2]-cycloaddition reactions [15,16]. The data obtained provide the convincing evidence that chemical behavior is governed to

a large extent by the stereochemical features of both chair and twist-boat structures, the bulkiness of substituents at C<sup>2</sup>, electronic ability of the latter and finally substrate solvation. Contrary to what is usually observed in chair-like six-membered saturated carbo- and heterocycles, flexible 2-R-1,3-dioxacyclohept-5-enes can serve as suitable substrates for thorough understanding of the conformer's reactivity [13]. Our long lasting interest in chemistry of cyclic acetals prompted us to study seven-membered fused acetals with incorporation of heteroaromatic planar fragments. Attention was drawn to the systems containing furan (**I**) and pyridine (**II**)—moieties (in the latter cases vitamin B<sub>6</sub> derivatives). As to furan containing molecules the choice of these systems was motivated by wide application of such heterocycles in synthetic chemistry [17–19].

\* Corresponding author. Fax: +7-8432-3872-78.

E-mail address: [evgenii.klimovitskii@ksu.ru](mailto:evgenii.klimovitskii@ksu.ru) (E.N. Klimovitskii).